

A SPLITTING CRITERION FOR RANK 2 VECTOR BUNDLES ON HYPERSURFACES IN $\mathbb{P}^4$

CARLO MADONNA

ABSTRACT. We show that Horrocks' criterion for the splitting of rank two vector bundles in $\mathbb{P}^3$ can be extended, with some assumptions on the Chern classes, on non singular hypersurfaces in $\mathbb{P}^4$. Extension of other splitting criterions are studied.

Introduction

A well known criterion of Horrocks ([OSS], pg.39) shows that a rank 2 vector bundle $\mathcal{E}$ on $\mathbb{P}^3$ splits as the direct sum of two line bundles if and only if $\oplus_n H^1(\mathbb{P}^3, \mathcal{E}(n)) = 0$. It is known that Horrocks criterion fails on a non singular hyperquadric of $\mathbb{P}^4$ (see [O]); moreover in [BGS, Remark 3) pg.169] it is remarked that, for any $r > 1$, there exists a non singular hypersurface $X \subset \mathbb{P}^4$ of degree r and a non splitting rank two vector bundle $\mathcal{E}$ on X with $\oplus_n H^1(\mathcal{E}(n)) = 0$, for $r > 2$, these bundles, form an infinite family.

One still can hope that, with a few number of exceptions on the numerical invariants of $\mathcal{E}$, also on a non singular hypersurface $X \subset \mathbb{P}^4$ of degree r , the vanishing $\oplus_n H^1(\mathcal{E}(n)) = 0$ forces a rank 2 vector bundle $\mathcal{E}$ on X to split.

Our main result is the following:

Theorem. *Let $X \subset \mathbb{P}^4$ be a non singular hypersurface, of degree r ; $\mathcal{E}$ be a rank 2 vector bundle on X with first Chern class c_1 ; let b be the maximum integer such that $h^0(\mathcal{E}(-b)) \neq 0$. Then: if $\oplus_n H^1(\mathcal{E}(n)) = 0$, $\mathcal{E}$ splits as the direct sum of two line bundles, unless $-r < 2b - c_1 < r - 2$.*

We remark that the number $-(2b - c_1)$ is a numerical invariant of $\mathcal{E}$, sometimes called "degree of stability": $2b - c_1 < 0$ is equivalent to $\mathcal{E}$ stable (in the sense of Mumford-Takemoto).

Examples show that on general hypersurfaces of degree up to 5, one side of the range is "sharp" and "almost sharp" the other one. Furthermore, we can show that for any $r > 5$ there are special smooth hypersurfaces of degree r for which one side of the bound is sharp. These bundles are obtained using an extended version of Serre correspondence between subcanonical curves and rank two vector bundles.

In section 3 we improve the criterion showing that for $-r < 2b - c_1 < r - 2$ condition $\oplus_n H^1(\mathcal{E}(n)) = 0$ may be replaced by the vanishing of $H^1(\mathcal{E}(n_0))$ for just one suitable $n_0 \in \mathbb{Z}$, extending in this way the Chiantini-Valabrega criterion (see [CV]).

We wish to thank Prof.L.Chiantini and Prof.S.Verra for their help in the preparation of this paper.

0. Notations and Conventions

With the word "variety" we shall indicate an irreducible projective algebraic variety defined over the complex field $\mathbb{C}$ while a rank two vector bundle is an algebraic rank 2 vector bundle. If $Y \subset X$ is a curve, $\mathcal{I}_{Y/X}$, or $\mathcal{I}_Y$ if there is no ambiguity, denote the ideal sheaf of Y in X and ω_Y its dualizing sheaf (see [H], ch.III, 7.5) If $\mathcal{F}$ is a sheaf on a variety X we define $h^i(\mathcal{F}) := \dim H^i(X, \mathcal{F})$.

When $\text{Pic}(X) \cong \mathbb{Z}$, we use this isomorphism to identify line bundles with integers. In particular, for any vector bundle $\mathcal{E}$ we consider $c_1(\mathcal{E}) = c_1 \in \mathbb{Z}$ and write $\mathcal{E}(n)$ for $\mathcal{E} \otimes \mathcal{O}_X(n)$. A locally complete intersection curve $Y \subset X$ is called a -subcanonical if $\omega_Y = \mathcal{O}_Y(a)$ for some $a \in \mathbb{Z}$.

We also define

$$b(\mathcal{E}) = b = \max\{n \mid h^0(\mathcal{E}(-n)) \neq 0\}.$$

We say that a rank 2 vector bundle $\mathcal{E}$ on X "splits" if it is isomorphic to the direct sum of two line bundles. We use the notion of stability as given in [OSS, pg. 160]. When $\text{Pic}(X) \cong \mathbb{Z}$ this means in our notation that a rank two vector bundle $\mathcal{E}$ is stable if and only if $2b - c_1 < 0$.

Note that the number $2b - c_1$ is invariant by twisting i.e

$$2b - c_1 = 2b(\mathcal{E}(n)) - c_1(\mathcal{E}(n))$$

$\forall n \in \mathbb{Z}$. If $b = 0$, as remarked in [H1, Remark 1.0.1], $\mathcal{E}$ has a global section whose zero-locus has codimension 2.

1. A well known criterion of Horrocks says that a rank 2 vector bundle $\mathcal{E}$ on $\mathbb{P}^3$ splits if and only if $\oplus_n H^1(\mathbb{P}^3, \mathcal{E}(n)) = 0$. In this section we obtain an extension of this criterion to rank 2 vector bundles on a non singular hypersurface $X \subset \mathbb{P}^4$.

Theorem 1.1. *Let $\mathcal{E}$ be a rank 2 vector bundle on X . Set $r = \deg(X)$. If $\oplus_n H^1(\mathcal{E}(n)) = 0$ then $\mathcal{E}$ splits unless $-r < 2b - c_1 < r - 2$.*

Proof. As remarked, the number $2b - c_1$ is an invariant of the bundle, so replacing $\mathcal{E}$ with $\mathcal{E}(-b)$, we can assume $b = 0$. We denote with H and L the intersection of X with a generic hyperplane and a generic plane in $\mathbb{P}^4$ and with $\mathcal{E}_H, \mathcal{E}_L$ the restrictions of $\mathcal{E}$ to H and L . By Bertini theorem there exist hyperplane sections so that H and L are non singular.

We split proof in two parts. First, we suppose that $-c_1 \geq r - 2$ and we prove theorem showing that:

- (1) if $H^1(\mathcal{E}(n)) = 0 \forall n \in \mathbb{Z}$ and $\mathcal{E}_H$ splits, then $\mathcal{E}$ splits;
- (2) if $H^1(\mathcal{E}_H(n)) = 0 \forall n \in \mathbb{Z}$ and $\mathcal{E}_L$ splits, then $\mathcal{E}_H$ splits;
- (3) $\mathcal{E}_L$ splits.

When $c_1 \geq r$ we prove something more, showing that either $h^1(\mathcal{E}(-2)) \neq 0$ or $h^1(\mathcal{E}(r - c_1 - 2)) \neq 0$.

Let us start with (1). Assume $\mathcal{E}_H = \mathcal{O}_H(\alpha) \oplus \mathcal{O}_H(\beta)$ with $\alpha \leq \beta$, for example. We have to extend this isomorphism to X . From the exact sequence

$$0 \rightarrow H^0(\mathcal{E}(n-1)) \rightarrow H^0(\mathcal{E}(n)) \rightarrow H^0(\mathcal{E}_H(n)) \rightarrow 0,$$

the pull back of the sections $s \in H^0(\mathcal{E}_H(-\alpha))$ and $s' \in H^0(\mathcal{E}_H(-\beta))$ gives two global sections $t \in H^0(\mathcal{E}(-\alpha))$ and $t' \in H^0(\mathcal{E}(-\beta))$, defining an injective morphism $\mu : \mathcal{O}_X(\alpha) \oplus \mathcal{O}_X(\beta) \rightarrow \mathcal{E}$ which is an isomorphism on H since t, t' are independent

on H . The locus where μ is not surjective is the zero-locus of the section $t \wedge t' \in H^0(\wedge^2 \mathcal{E}(-\alpha - \beta))$, which is empty or has codimension one, since $\wedge^2 \mathcal{E}(-\alpha - \beta)$ is of rank 1. This locus is necessary empty, otherwise it would intersect H while the sections t, t' are independent on H . So t, t' are independent on X and $\mathcal{E}$ splits.

The same argument apply to prove (2).

To prove (3), note that since we are assuming $b(\mathcal{E}) = 0$, $\mathcal{E}$ has a global section whose zero-locus has codimension two, say $Y = (s)_0$. Hence $\mathcal{E}_L$ has a nowhere vanishing global section, and this one define an injective morphism $s : \mathcal{O}_L \rightarrow \mathcal{E}_L$, where $\mathcal{F} = \mathcal{E}_L / \text{Im}(s)$ is a subbundle of $\mathcal{E}_L$. By the exact sequence

$$0 \rightarrow \mathcal{O}_L \rightarrow \mathcal{E}_L \rightarrow \mathcal{F} \rightarrow 0 \quad (*)$$

we have $\text{rk}(\mathcal{F}) = 1$ and comparing Chern classes, we find $\mathcal{F} = \mathcal{O}_L(c_1)$. Then (*) reads as:

$$0 \rightarrow \mathcal{O}_L \rightarrow \mathcal{E}_L \rightarrow \mathcal{O}_L(c_1) \rightarrow 0,$$

hence $\mathcal{E}_L$ is an extension of $\mathcal{O}_L(c_1)$ by $\mathcal{O}_L$ i.e an element of the group

$$\text{Ext}^1(\mathcal{O}_L(c_1), \mathcal{O}_L) \cong H^1(\mathcal{O}_L(-c_1)).$$

Using $c_1 + r - 3 < 0$ we have $h^1(\mathcal{O}_L(-c_1)) = h^0(\mathcal{O}_L(c_1 + r - 3)) = 0$ by duality, since L is a plane curve of degree r ; so the only possible extension is the trivial one, i.e $\mathcal{E}_L = \mathcal{O}_L \oplus \mathcal{O}_L(c_1)$.

This proves (3), so one half of the theorem is proved.

Assume now that $c_1 \geq r$. In this case $\mathcal{E}$ is stable and so it does not split. Indeed if $\mathcal{E} = \mathcal{O}_X(\alpha) \oplus \mathcal{O}_X(\beta)$, being $c_1 = \alpha + \beta$, it is $\alpha > 0$ or $\beta > 0$ and so $h^0(\mathcal{E}(-1)) > 0$, absurd since $b = 0$. We will prove that in this case $h^1(\mathcal{E}(-2)) \neq 0$ or $h^1(\mathcal{E}(-c_1 + r - 2)) \neq 0$. From the exact sequence

$$0 \rightarrow \mathcal{E}(-2) \rightarrow \mathcal{E}(-1) \rightarrow \mathcal{E}_H(-1) \rightarrow 0$$

one gets:

$$\dots \rightarrow H^0(\mathcal{E}(-1)) \rightarrow H^0(\mathcal{E}_H(-1)) \rightarrow H^1(\mathcal{E}(-2)) \rightarrow \dots,$$

hence if $h^0(\mathcal{E}_H(-1)) \neq 0$ then $h^1(\mathcal{E}(-2)) \neq 0$ too, since $h^0(\mathcal{E}(-1)) = 0$. Assume $h^1(\mathcal{E}(-2)) = 0$. We will show that $h^1(\mathcal{E}(-c_1 + r - 2)) = 0$ implies $h^0(\mathcal{E}_H(-1)) \neq 0$, proving in this way that if $h^1(\mathcal{E}(-2)) = 0$ then $h^1(\mathcal{E}(-c_1 + r - 2)) \neq 0$. Indeed assume $h^0(\mathcal{E}_H(-1)) = 0$; from

$$0 \rightarrow \mathcal{E}_H(-2) \rightarrow \mathcal{E}_H(-1) \rightarrow \mathcal{E}_L(-1) \rightarrow 0$$

we have an exact sequence

$$\dots \rightarrow H^0(\mathcal{E}_H(-1)) \rightarrow H^0(\mathcal{E}_L(-1)) \rightarrow H^1(\mathcal{E}_H(-2)) \rightarrow \dots$$

Suppose we have proved that $h^0(\mathcal{E}_L(-1)) \neq 0$. Then $h^1(\mathcal{E}_H(-2)) \neq 0$ and the sequence

$$\dots \rightarrow H^1(\mathcal{E}(-2)) \rightarrow H^1(\mathcal{E}_H(-2)) \rightarrow H^2(\mathcal{E}(-3)) \rightarrow \dots$$

proves that $h^1(\mathcal{E}(-2)) \neq 0$ or $h^2(\mathcal{E}(-3)) \neq 0$, absurd since by Serre duality $h^2(\mathcal{E}(-3)) = h^1(\mathcal{E}(-c_1 + r - 2)) = 0$ and by assumption $h^1(\mathcal{E}(-2)) = 0$.

It remains to show that $h^0(\mathcal{E}_L(-1)) \neq 0$. If not, by the exact sequence

$$0 \rightarrow \mathcal{O}_L(-1) \rightarrow \mathcal{E}_L(-1) \rightarrow \mathcal{O}_L(c_1 - 1) \rightarrow 0$$

we have an injective map

$$H^0(\mathcal{O}_L(c_1 - 1)) \rightarrow H^1(\mathcal{O}_L(-1)) \cong (H^0(\mathcal{O}_L(r - 2)))^\vee$$

absurd since $c_1 - 1 \geq r - 1 \geq 0$. $\square$

By the previous theorem we have the following:

Corollary 1.2. *If $\mathcal{E}$ is a non splitting rank 2 vector bundle on X and $2b - c_1 \geq r - 2$ then there is an integer n_0 so that $h^1(\mathcal{E}(n_0)) \neq 0$.*

From the proof of the theorem, it follows:

Corollary 1.3. *Let $\mathcal{E}$ be a rank 2 vector bundle on X and $2b - c_1 \leq -r$. Then $h^1(\mathcal{E}(-b - 2)) \neq 0$ or $h^1(\mathcal{E}(-c_1 + b + r - 2)) \neq 0$.*

Remark. If $r = 1$ i.e $X = \mathbb{P}^3$ theorem 1.1 gives Horrocks splitting criterion.

2. In this section we will give examples of non splitting rank 2 vector bundles with $\oplus_n H^1(\mathcal{E}(n)) = 0$ showing that one side of the range of Theorem 1.1 is “sharp”. We will use an extended version of Serre correspondence between curves and bundles. The same proof of [H1] or [GH] works, under the hypothesis $h^1(\mathcal{L}^\vee) = h^2(\mathcal{L}^\vee) = 0$, which is always fulfilled when X is a smooth hypersurface in $\mathbb{P}^4$ and $\mathcal{L}$ is a line bundle on X (for $\text{Pic}(X) \cong \mathbb{Z}$, see [H1, Remark 1.1.1]), in this case too. First a definition:

Definition. Let $\mathcal{E}$ be a rank 2 vector bundle on a non singular 3-dimensional variety, $s \in H^0(\mathcal{E})$ a global section whose zero-locus $C = (s)_0$ has codimension 2. Let $\mathcal{L} \in \text{Pic}(X)$ with $h^1(\mathcal{L}^\vee) = h^2(\mathcal{L}^\vee) = 0$ and let $\phi : \wedge^2 \mathcal{E} \rightarrow \mathcal{L}$ be an isomorphism. We say that two triples $(\mathcal{E}, s, \phi)$ and $(\mathcal{E}', s', \phi')$ are isomorphic if there is an isomorphism $\psi : \mathcal{E} \rightarrow \mathcal{E}'$ and an element $\lambda \in \mathbb{C}^*$, such that $s' = \lambda\psi(s)$ and $\phi' = \lambda^2\phi(\wedge^2\psi)^{-1}$.

The correspondence between bundles and curves is given by the following:

Theorem 2.1(Serre). *Let X be a non singular 3-dimensional variety. A curve $C \subset X$ occurs as the zero-locus of a section of a rank 2 vector bundle $\mathcal{E}$ on X if it is a local complete intersection and ω_C is isomorphic to the restriction to C of some invertible sheaf $\mathcal{L}$ on X with $h^1(\mathcal{L}^\vee) = h^2(\mathcal{L}^\vee) = 0$. More precisely, for any fixed invertible sheaf $\mathcal{L}$ on X with the previous conditions, there exist a bijection between the following set of data:*

- (i) *the set of triples $(\mathcal{E}, s, \phi)$ modulo isomorphism;*
- (ii) *the set of pairs (C, ξ) where C is a locally complete intersection curve in X , and $\xi : \mathcal{L} \otimes \omega_X \otimes \mathcal{O}_C \rightarrow \omega_C$ is an isomorphism (modulo $\mathbb{C}^*$).*

From the theorem it follows:

Corollary 2.2. *Let $C \subset X$ be a locally complete intersection, subcanonical curve and $\mathcal{E}$ the bundle associated to C . Then C is complete intersection if and only if $\mathcal{E}$ splits.*

Proposition 2.3. *Let X be a non singular 3-dimensional variety and $\mathcal{E}$ a rank 2 vector bundle on X . If $\mathcal{E}$ is generated by global sections, then the zero-locus of a general section is non singular.*

Proof. The same proof of [HM, Theorem 5.1] works in this case too. $\square$

Remark.

- (1) If C is reduced and connected the bundle $\mathcal{E}$ associated to C is unique;
- (2) if $n \gg 0$ then $\mathcal{E}(n)$ is generated by global sections and if $s \in H^0(\mathcal{E}(n))$ is general then $(s)_0$ is a non singular irreducible curve;

- (3) if $X \subset \mathbb{P}^4$ is a non singular hypersurface and $\mathcal{E}$ is associated to a curve $C \subset X$ with $\deg(C) = d$ and genus g then $d = c_2(\mathcal{E})$, $2g - 2 = c_2(c_1 + r - 5)$ and C is $(c_1 + r - 5)$ -subcanonical.

Remark. Let $C \subset \mathbb{P}^4$ be a curve, complete intersection of three non singular hypersurfaces of degree l, m, n , contained in a non singular hypersurface $X \subset \mathbb{P}^4$. Assume that C is not complete intersection in X and let $\mathcal{E}$ associated to C on X . Then $\oplus_n H^1(\mathcal{E}(n)) = 0$. Indeed using the exact sequences

$$0 \rightarrow \mathcal{I}_{X/\mathbb{P}^4} \rightarrow \mathcal{I}_{C/\mathbb{P}^4} \rightarrow \mathcal{I}_{C/X} \rightarrow 0$$

$$0 \rightarrow \mathcal{O}_X \rightarrow \mathcal{E} \rightarrow \mathcal{I}_C(c_1) \rightarrow 0$$

we have,

$$\dots \rightarrow H^1(\mathcal{I}_{C/\mathbb{P}^4}(n)) \rightarrow H^1(\mathcal{I}_{C/X}(n)) \rightarrow H^2(\mathcal{I}_{X/\mathbb{P}^4}(n)) \rightarrow \dots$$

so that $H^1(\mathcal{I}_{C/X}(n)) = 0$, hence from the second sequence we have $H^1(\mathcal{E}(n)) = 0$ for all n .

We call a curve obtained in this way a "curve of type (l, m, n) ".

So if $C \subset \mathbb{P}^4$ is a curve of type (l, m, n) , contained in a non singular hypersurface $X \subset \mathbb{P}^4$ with $\deg(X) = r > n \geq m \geq l > 0$, then C is $(l + m + n - 5)$ -subcanonical, and if $\mathcal{E}$ is associated to C we have:

- (1) $c_1 = l + n + m - r$ and $c_2 = \deg(C) = lmn$;
- (2) $\oplus_n H^1(\mathcal{E}(n)) = 0$.

Let $C = (l, m, n)$ as above with $0 < l \leq m \leq n < r$. If $m + n \leq r$ then $c_1 = l + m + n - r \leq l$, since $m + n \leq r$, hence $b = 0$ and $2b - c_1 = r - l - m - n$.

If $m + n \geq r$ then $c_1 = l + m + n - r \geq l$ and $b = m + n - r$, hence $2b - c_1 = m + n - r - l$ since $m - l \geq 0$ and $n - r \geq -\frac{r}{2}$.

Observe that in both cases, we always have $2b - c_1 \geq -\frac{r}{2}$.

Example 2.4. Let $X \subset \mathbb{P}^4$ be a general quintic hypersurface, and let $C \subset X$ be a line, so C is (-2) -subcanonical. As remarked we have a bundle $\mathcal{E}$ on X associated to C with $\oplus_n H^1(\mathcal{E}(n)) = 0$; this shows sharpness on one side, i.e when $2b - c_1 = 2$, for **all quintics**. Indeed by $0 \rightarrow H^0(\mathcal{O}_X) \rightarrow H^0(\mathcal{E}) \rightarrow H^0(\mathcal{I}_C(-2)) \rightarrow 0$ we find $b = 0$ hence $2b - c_1 = 2$. Note that $\mathcal{E}$ does not split since C is not complete intersection in X .

Example 2.5. Let $C \subset \mathbb{P}^4$ a line. By Bertini there exist a non singular hypersurface X containing C , of any degree $r > 0$. If $\mathcal{E}$ is associated to C , as remarked, we have $\oplus_n H^1(\mathcal{E}(n)) = 0$ and $2b - c_1 = r - 3$. So the right bound of Theorem 1.1 is sharp on hypersurfaces containing a line.

If $\deg(X) = 2$, the range of Theorem 1.1 is sharp. Indeed the only exception is for $2b - c_1 = -1$ and in fact we have bundles associated to lines, which do not split.

Example 2.6. For $2b - c_1$ negative, the best we can do is to consider curves of type $(\frac{r}{2}, \frac{r}{2}, \frac{r}{2})$ for r even, or $(\frac{r-1}{2}, \frac{r-1}{2}, \frac{r+1}{2})$ for $r > 1$ odd. From the Remark, we know that there exists a smooth hypersurface of degree r containing C ; the corresponding rank 2 bundle $\mathcal{E}$ has no cohomology in the middle and it does not split. As remarked, we have $2b - c_1 = r - l - m - n = -\frac{r}{2}$ for r even or $2b - c_1 = -\frac{r-1}{2}$ for $r > 1$ odd.

Observe that even if these examples are far from our bound yet they give stable rank 2 bundles with no cohomology in the middle (but only on some particular hypersurface).

3. In the present section we weaken the criterion proved above, showing that condition $\oplus_n H^1(\mathcal{E}(n)) = 0$ may be replaced by the vanishing of some $H^1(\mathcal{E}(n_0))$ for just one suitable n_0 , extending in this way the Chiantini-Valabrega criterion. The idea is the same used in [CV] i.e we find an integer n_0 such that:

- (1) if $n \geq n_0$ and $H^1(\mathcal{E}(n)) = 0$ then $H^1(\mathcal{E}(n+1)) = 0$;
- (2) if $n \leq n_0$ and $H^1(\mathcal{E}(n)) = 0$ then $H^1(\mathcal{E}(n-1)) = 0$.

In this section we always twist $\mathcal{E}$ suitably, so that we may assume $b(\mathcal{E}) = 0$. We have to prove a preliminary vanishing:

Proposition 3.1. *Let $m > r - 4$, $H^1(\mathcal{E}(m)) = 0$ and $H^1(\mathcal{E}(-m - c_1 + r - 4)) = 0$. Then $H^1(\mathcal{E}(m+1)) = 0$.*

Proof. By the exact sequence:

$$0 \rightarrow \mathcal{E}(m-1) \rightarrow \mathcal{E}(m) \rightarrow \mathcal{E}_H(m) \rightarrow 0$$

and by Serre duality we have the following exact sequence

$$\dots \rightarrow H^1(\mathcal{E}(m)) \rightarrow H^1(\mathcal{E}_H(m)) \rightarrow H^2(\mathcal{E}(m-1)) \cong H^1(\mathcal{E}(-m - c_1 + r - 4))^\vee \rightarrow \dots$$

which, by assumption, gives $h^1(\mathcal{E}_H(m)) = 0$. By the exact sequence:

$$\dots \rightarrow H^1(\mathcal{E}(m)) \rightarrow H^1(\mathcal{E}(m+1)) \rightarrow H^1(\mathcal{E}_H(m+1)) \rightarrow \dots$$

using $h^1(\mathcal{E}(m)) = 0$ it is sufficient to show that $h^1(\mathcal{E}_H(m+1)) = 0$. By

$$0 \rightarrow \mathcal{E}_H(m) \rightarrow \mathcal{E}_H(m+1) \rightarrow \mathcal{E}_L(m+1) \rightarrow 0$$

$$\dots \rightarrow H^1(\mathcal{E}_H(m)) \rightarrow H^1(\mathcal{E}_H(m+1)) \rightarrow H^1(\mathcal{E}_L(m+1)) \xrightarrow{\alpha} H^2(\mathcal{E}_H(m)) \rightarrow \dots$$

since $h^1(\mathcal{E}_H(m)) = 0$, it is sufficient to show that α is injective i.e that the dual map

$$\alpha^\vee : H^0(\mathcal{E}_H(-m - c_1 + r - 4)) \rightarrow H^0(\mathcal{E}_L(-m - c_1 + r - 4))$$

is surjective. To prove surjectivity of $\alpha^\vee$ remind that $b = 0$ and $\text{rk}(\mathcal{E}) = 2$, hence $\mathcal{E}$ has a global section whose zero-locus has codimension two, and so for a generic L , $\mathcal{E}_L$ has a nowhere vanishing global section giving the following exact sequence:

$$0 \rightarrow \mathcal{O}_L \rightarrow \mathcal{E}_L \rightarrow \mathcal{O}_L(c_1) \rightarrow 0.$$

From this one and by the commutativity of the following diagramm:

$$\begin{array}{ccccccc} 0 & \longrightarrow & H^0 \mathcal{O}_H(-m - c_1 + r - 4) & \longrightarrow & H^0 \mathcal{E}_H(-m - c_1 + r - 4) & & \\ & & \beta \downarrow & & \alpha^\vee \downarrow & & \\ 0 & \longrightarrow & H^0 \mathcal{O}_L(-m - c_1 + r - 4) & \xrightarrow{\gamma} & H^0 \mathcal{E}_L(-m - c_1 + r - 4) & \longrightarrow & 0 \\ & & \downarrow & & & & \\ & & 0 & & & & \end{array}$$

since γ is bijective being $h^0 \mathcal{O}_L(-m + r - 4) = 0$ and β is surjective too, since L is a plane curve hence arithmetically normal, then $\alpha^\vee$ is surjective. $\square$

Another preliminary vanishing result is the following:

Proposition 3.2. *Let $m \leq -c_1$. If $H^1(\mathcal{E}(m)) = 0$ then $H^1(\mathcal{E}(m-1)) = 0$.*

Proof. By the exact sequence:

$$\dots \rightarrow H^0(\mathcal{E}(m)) \xrightarrow{u} H^0(\mathcal{E}_H(m)) \rightarrow H^1(\mathcal{E}(m-1)) \rightarrow \dots$$

if u surjects then $h^1(\mathcal{E}(m-1)) = 0$. We assume u not surjective and we will show that $\mathcal{E}$ splits. For $n \gg 0$, as remarked, $\mathcal{E}(n)$ has a global section whose zero-locus is a irreducible non singular curve Y and

$$c_2(\mathcal{E}(n)) = \deg(C) = c_2 + rnc_1 + rn^2.$$

From Koszul complex of this section we have an exact sequence

$$0 \rightarrow \mathcal{O}_X \rightarrow \mathcal{E}(n) \rightarrow \mathcal{I}_C(c_1 + 2n) \rightarrow 0,$$

using $h^0(\mathcal{E}(-n)) = 0 \ \forall n > 0$ and $h^0\mathcal{E} \neq 0$, we find $h^0(\mathcal{I}_C(c_1 + n)) \neq 0$ hence there is a non zero sections $s \in H^0(\mathcal{E})$, giving a divisor S on X of type $(c_1 + n)H$ containing C , and $c_1 + n$ is the minimum integer α such that $C \subset \alpha H$. Moreover S is irreducible, indeed if not C should be contained in some component of S , contradicting the minimality of $c_1 + n$. Let h be a general hyperplane divisor on H and set $\Gamma := C \cap h \subset S \cap h := C''$. Note that C'' is irreducible by Bertini theorem since S is irreducible. Next, if $h^0(\mathcal{E}_H(m)) \neq 0$, $\mathcal{E}_H(m)$ has a global section independent from s . By the exact sequence:

$$0 \rightarrow \mathcal{O}_H \rightarrow \mathcal{E}_H(n) \rightarrow \mathcal{I}_\Gamma(c_1 + 2n) \rightarrow 0 \quad (*)$$

Γ is also contained in a divisor C' on H , of type $(c_1 + n + m)h$ independent from C' . Then Γ is contained in C'' and in C' and so by intersection formula, we find

$$\deg(\Gamma) \leq (c_1 + n)h \cdot (c_1 + n + m)h = r(c_1 + n)(c_1 + n + m) \leq r(c_1 + n)n$$

since $m + c_1 \leq 0$. Then

$$c_2 + rnc_1 + rn^2 \leq r(c_1 + n)n = rnc_1 + rn^2$$

i.e $c_2 \leq 0$. Being $\deg(Y) = c_2 \geq 0$ it is $c_2 = 0$ and $Y = \emptyset$. So $\mathcal{E}(n)$ has a never vanishing global section i.e $\mathcal{E}(n)$ splits, proving theorem. $\square$

We can now prove the splitting criterion for unstable rank 2 vector bundles.

Theorem 3.3. *Let $\mathcal{E}$ be a non splitting rank 2 vector bundle and $c_1 \leq 2 - r$.*

- (1) *If $r + c_1$ is odd then $H^1(\mathcal{E}(\frac{-c_1+r-3}{2})) \neq 0$;*
- (2) *if $r + c_1$ is even then $H^1(\mathcal{E}(\frac{-c_1+r-4}{2})) \neq 0$ and $H^1(\mathcal{E}(\frac{-c_1+r-2}{2})) \neq 0$. Moreover if $c_1 \leq -r$ then $H^1(\mathcal{E}(\frac{-c_1+r}{2})) \neq 0$ too.*

Proof. (2) Assume $h^1(\mathcal{E}(\frac{-c_1+r-4}{2})) = 0$. By Proposition 3.2 we find $h^1(\mathcal{E}(n)) = 0$ for all $n \leq \frac{-c_1+r-4}{2}$. Setting $m = \frac{-c_1+r-4}{2}$ we have

$$h^1(\mathcal{E}(-m - c_1 + r - 4)) = h^1(\mathcal{E}(\frac{-c_1 + r - 4}{2})) = 0$$

and $m > r - 4$ so by Proposition 3.1 we have $h^1(\mathcal{E}(n)) = 0$ for all $n \geq \frac{-c_1+r-4}{2}$. It follows that $h^1\mathcal{E}(n) = 0 \ \forall n$ and $\mathcal{E}$ splits by Theorem 1.1, absurd. The same argument apply to prove the remaining non vanishing with the extra hypothesis $c_1 \leq -r$ required for $h^1(\mathcal{E}(\frac{-c_1+r}{2})) = 0$.

- (1) Similar proof works in this case too. $\square$

Now let us turn our attention to the range $c_1 \geq r$. We are able to prove:

Proposition 3.4. *Let $m \geq 0$. If $H^1(\mathcal{E}(-m-2)) = 0$ then $H^1(\mathcal{E}(n)) = 0$ for all $n \leq -m-2$.*

Proof. By the exact sequence

$$\dots \rightarrow H^0(\mathcal{E}(-m-2)) \rightarrow H^0(\mathcal{E}_H(-m-2)) \rightarrow H^1(\mathcal{E}(-m-3)) \rightarrow 0,$$

we find $H^1(\mathcal{E}(-m-3)) \cong H^0(\mathcal{E}_H(-m-2))$ since $h^0(\mathcal{E}(-m-2)) = 0$ being $-m-2 < -1$. By the exact sequence

$$\dots \rightarrow H^0(\mathcal{E}(-m-1)) \rightarrow H^0(\mathcal{E}_H(-m-1)) \rightarrow H^1(\mathcal{E}(-m-2)) \rightarrow \dots,$$

since $h^0(\mathcal{E}(-m-1)) = h^1(\mathcal{E}(-m-2)) = 0$ we find $h^0(\mathcal{E}_H(-m-2)) = 0$, following thesis. $\square$

Applying previous theorem to stable bundles we have the following:

Corollary 3.5. *Let $c_1 \geq r$. Then $H^1(\mathcal{E}(-2)) \neq 0$.*

Proof. If $h^1(\mathcal{E}(-2)) = 0$, setting $m = 0$ by previous proposition, we find $h^1(\mathcal{E}(-c_1 + r - 2)) = 0$ since $-c_1 + r - 2 \leq -2$, absurd by Corollary 1.3. $\square$

Corollary 3.6. *Let $c_1 \geq r$.*

- (1) *If $c_1 + r$ is odd and $H^1(\mathcal{E}(\frac{-c_1+r-3}{2})) = 0$ then $H^1(\mathcal{E}(n)) = 0$ for all $n \leq \frac{-c_1+r-3}{2}$;*
- (2) *if $c_1 + r$ is even and $H^1(\mathcal{E}(\frac{-c_1+r-4}{2})) = 0$ then $H^1(\mathcal{E}(n)) = 0$ for all $n \leq \frac{-c_1+r-4}{2}$.*

Proof. (1) Setting $m = \frac{c_1-r-1}{2}$ hypothesis reads as

$$h^1(\mathcal{E}(-m-2)) = h^1(\mathcal{E}(\frac{-c_1+r-3}{2})) = 0.$$

Thesis follows by Proposition 3.4.

(2) The same argument of (1) applies. $\square$

Remark. If moreover $c_1 \geq r + 1$ we can prove in the same way that:

if $H^1(\mathcal{E}(\frac{-c_1+r-2}{2})) = 0$ then $H^1(\mathcal{E}(n)) = 0$ for all $n \leq \frac{-c_1+r-2}{2}$ while if $c_1 \geq r + 3$ and $H^1(\mathcal{E}(\frac{-c_1+r}{2})) = 0$ then $H^1(\mathcal{E}(n)) = 0$ for all $n \leq \frac{-c_1+r}{2}$.

In the other side, we have:

Proposition 3.7. *Let $c_1 \geq r$.*

- (1) *If $c_1 + r$ is odd and $H^1(\mathcal{E}(\frac{-c_1+r-3}{2})) = 0$ then $H^1(\mathcal{E}(n)) = 0$ for all $n \geq \frac{-c_1+r-3}{2}$;*
- (2) *if $c_1 + r$ is even and $H^1(\mathcal{E}(\frac{-c_1+r-4}{2})) = 0$ then $H^1(\mathcal{E}(n)) = 0$ for all $n \geq \frac{-c_1+r-4}{2}$.*

Proof. (1) For $m \gg 0$, as remarked, $\mathcal{E}(m)$ has a global section whose zero-locus is an irreducible non singular curve C giving the following exact sequence

$$0 \rightarrow \mathcal{O}_X \rightarrow \mathcal{E}(m) \rightarrow \mathcal{I}_C(c_1 + 2m) \rightarrow 0. \quad (*)$$

By hypothesis

$$h^1(\mathcal{E}(\frac{-c_1+r-3}{2})) = h^1(\mathcal{J}_C(\frac{c_1+r-3}{2} + m)) = 0,$$

and being $\frac{-c_1+r-3}{2} < 0$ we have

$$h^0(\mathcal{E}(\frac{-c_1+r-3}{2})) = h^0(\mathcal{J}_C(\frac{c_1+r-3}{2} + m)) = 0$$

too, hence by the exact sequence

$$0 \rightarrow \mathcal{J}_C(\frac{c_1+r-3}{2} + m) \rightarrow \mathcal{O}_X(\frac{c_1+r-3}{2} + m) \rightarrow \mathcal{O}_C(\frac{c_1+r-3}{2} + m) \rightarrow 0$$

we find

$$h^0(\mathcal{O}_X(\frac{c_1+r-3}{2} + m)) = h^0(\mathcal{O}_C(\frac{c_1+r-3}{2} + m)).$$

By Riemann-Roch, being C a locally complete intersection and $(c_1 + 2m + r - 5)$ -subcanonical curve, we see that

$$h^0(\mathcal{O}_C(\frac{c_1+r-3}{2} + m)) - h^0(\mathcal{O}_C(\frac{c_1+r-7}{2} + m)) = d. \quad (\text{I})$$

Note that

$$h^0(\mathcal{O}_X(\frac{c_1+r-7}{2} + m)) = h^0(\mathcal{O}_C(\frac{c_1+r-7}{2} + m)),$$

indeed, $h^0(\mathcal{E}(\frac{-c_1+r-7}{2})) = 0$ since $\frac{-c_1+r-7}{2} < -3$ and by Corollary 3.6, we have

$$h^1(\mathcal{E}(\frac{-c_1+r-7}{2})) = h^1(\mathcal{J}_C(\frac{c_1+r-7}{2} + m)) = 0$$

so previous equality holds and (I) reads as:

$$d = h^0(\mathcal{O}_X(\frac{c_1+r-3}{2} + m)) - h^0(\mathcal{O}_X(\frac{c_1+r-7}{2} + m)). \quad (\text{II})$$

Now assume $h^1(\mathcal{E}(\frac{-c_1+r-1}{2})) \neq 0$. From (*) we have $h^1(\mathcal{J}_C(\frac{c_1+r-1}{2} + m)) \neq 0$ too, hence:

$$h^0(\mathcal{O}_C(\frac{c_1+r-1}{2} + m)) > h^0(\mathcal{O}_X(\frac{c_1+r-1}{2} + m)).$$

Again using Riemann-Roch we find

$$h^0(\mathcal{O}_C(\frac{c_1+r-1}{2} + m)) - h^0(\mathcal{O}_C(\frac{c_1+r-9}{2} + m)) = 2d. \quad (\text{III})$$

Note that

$$h^0(\mathcal{O}_X(\frac{c_1+r-9}{2} + m)) = h^0(\mathcal{O}_C(\frac{c_1+r-9}{2} + m)).$$

Indeed we have $h^0(\mathcal{J}_C(\frac{c_1+r-9}{2} + m)) = 0$ since $\frac{-c_1+r-9}{2} < \frac{-c_1+r-7}{2} < 0$, and $h^1(\mathcal{J}_C(\frac{c_1+r-9}{2} + m)) = 0$ by Corollary 2.6, since $\frac{-c_1+r-9}{2} < \frac{c_1+r-3}{2}$. Hence (III) is given by:

$$2d > h^0(\mathcal{O}_X(\frac{c_1+r-1}{2} + m)) - h^0(\mathcal{O}_X(\frac{c_1+r-9}{2} + m)). \quad (\text{IV})$$

Comparing (II) and (IV) we find

$$\begin{aligned} 2[h^0(\mathcal{O}_X(\frac{c_1+r-3}{2}+m)) - h^0(\mathcal{O}_X(\frac{c_1+r-7}{2}+m))] &> \\ &> h^0(\mathcal{O}_X(\frac{c_1+r-1}{2}+m)) - h^0(\mathcal{O}_X(\frac{c_1+r-9}{2}+m)). \end{aligned} \quad (V)$$

By standard calculus we find

$$\begin{aligned} h^0(\mathcal{O}_X(k)) - h^0(\mathcal{O}_X(p)) &= \frac{1}{4!}[4r(k^3 - p^3) + 6r(5-r)(k^2 - p^2) + \\ &\quad + 2r(2r^2 - 15r + 35)(k - p)] \end{aligned}$$

hence (V) gives the following:

$$2[-\frac{r}{12} + \frac{c_1^2 r}{4} + c_1 m r + m^2 r + \frac{r^3}{12}] > \frac{11r}{6} + \frac{c_1^2 r}{2} + 2c_1 m r + 2m^2 r + \frac{r^3}{6}$$

and one realizes soon, by standard calculus, that this implies $r < 0$, absurd.

(2) The same argument of (1) applies. $\square$

Remark. In the same way we can prove that:

if $c_1 + r$ is even, $c_1 \geq r$ and $H^1(\mathcal{E}(\frac{-c_1+r-2}{2})) = 0$ then $H^1(\mathcal{E}(n)) = 0 \quad \forall n \geq \frac{-c_1+r-2}{2}$. Moreover if $c_1 \geq r + 2$ and $H^1(\mathcal{E}(\frac{-c_1+r}{2})) = 0$ then $H^1(\mathcal{E}(n)) = 0 \quad \forall n \geq \frac{-c_1+r}{2}$.

From Theorem 3.3, Corollary 3.6 and Proposition 3.7 we have our main splitting criterion:

Theorem 3.8. *Let $\mathcal{E}$ be a rank 2 vector bundle on X . Then:*

- (1) *if $c_1 + r$ is odd, $\mathcal{E}$ splits if and only if $H^1(\mathcal{E}(\frac{-c_1+r-3}{2})) = 0$;*
- (2) *if $c_1 + r$ is even, $\mathcal{E}$ splits if and only if $H^1(\mathcal{E}(\frac{-c_1+r-4}{2})) = 0$,*

unless $-r < -c_1 < r - 2$.

Remark. If $c_1 \leq 2 - r$ or $c_1 \geq r + 2$ we find another suitable vanishing: $\mathcal{E}$ splits if and only if $H^1(\mathcal{E}(\frac{-c_1+r-2}{2})) = 0$. Moreover if $c_1 \leq -r$ or $c_1 \geq r + 4$ then $\mathcal{E}$ splits if and only if $H^1(\mathcal{E}(\frac{-c_1+r}{2})) = 0$.

Note that for $r = 1$ i.e if $X = \mathbb{P}^3$ we obtain the Chiantini-Valabrega splitting criterion ([CV]), while for $r = 2$ we get a result of Ottaviani ([O, Theorem 3.8]).

Remark. Results presented in this paper concern obviously, via Serre correspondence, complete intersection and projective normality. Indeed by the exact sequences

$$0 \rightarrow \mathcal{O}_X \rightarrow \mathcal{E} \rightarrow \mathcal{I}_C(c_1) \rightarrow 0$$

if C is a -subcanonical then $\omega_C = \mathcal{O}_C(a) = \mathcal{O}_C(c_1 + r - 5)$ so previous theorem reads as follow: C is complete intersection if and only if:

- (1) $H^1(\mathcal{I}_C(\frac{a}{2} + 1)) = 0$, if a is even;
- (2) $H^1(\mathcal{I}_C(\frac{a+1}{2})) = 0$, if a is odd,

unless $-3 < a < 2r - 5$.

The following results are a natural generalization of the previous ones and can be proved by a straightforward arrangement of the previous procedure.

Theorem 3.9. *Let X be a non singular 3-dimensional variety. Assume $\text{Pic}(X) \cong \mathbb{Z}$, $h^1(\mathcal{O}_X(n)) = 0$ for all $n \in \mathbb{Z}$. If $\omega_X = \mathcal{O}_X(e)$ then: if $\oplus_n H^1(\mathcal{E}(n)) = 0$ then $\mathcal{E}$ splits unless $-e - 5 < 2b - c_1 < e + 3$.*

Theorem 3.10. *Let X be as above and $\mathcal{E}$ a rank 2 vector bundle on X . Then $\mathcal{E}$ splits if and only if:*

- (1) $H^1(\mathcal{E}(\frac{-c_1+e+2}{2})) = 0$, if $c_1 + e$ is even;
- (2) $H^1(\mathcal{E}(\frac{-c_1+e+1}{2})) = 0$, if $c_1 + e$ is odd.

unless $-e - 5 < 2b - c_1 < e + 3$.

REFERENCES

- [CV] L.Chiantini-P.Valabrega, *Subcanonical curves and complete intersections in projective 3-Space*, Ann. Mat. Pura Appl. (4) **138** (1984), 309–330.
- [BGS] R.O.Buchweitz-G.M.Greuel-F.O.Schreyer, *Cohen-Macaulay modules on hypersurface singularities II*, Invent. Math. **88** (1987), 165–182.
- [GH] P.Griffiths-J.Harris, *Two proofs of a theorem concerning algebraic space curve*, Proceeding of the Eighth National Mathematics Conference (Arya-Mehr Univ. Tech., Teheran, 1977), pp. 350–370.
- [H] R.Hartshorne, *Algebraic Geometry* (GTM 52) (Springer-Verlag, eds.), Berlin, 1993.
- [H1] ———, *Stable vector bundles of rank 2 on $\mathbb{P}^3$* , Math. Ann. **238** (1978), 229–280.
- [HM] G.Horrocks-D.Mumford, *A rank 2 vector bundle on $\mathbb{P}^4$ with 15,000 symmetries*, Topology **12** (1973), 63–81.
- [OSS] C.Okonek-M.Schneider-H.Spindler, *Vector Bundles on Complex Projective Spaces* (Prog. Math. 3) (Birkhäuser, eds.), Boston, Mass., 1980.
- [O] G.Ottaviani, *Some extension of Horrocks criterion to vector bundles on Grassmannians and quadrics*, Ann. Mat. Pura Appl. (4) **155** (1989), 317–341.

MADONNA CARLO, Pz.ACCADEMIA ANTIQUARIA 30, 00147 ROMA
E-mail address: `madonna@ciop.mat.uniroma3.it`